

AJOVY[®]
(fremanezumab-vfrm)
injection 225 mg/1.5 mL



AJOVY COULD GIVE YOU THE CHANCE TO BE MIGRAINE FREE

NOW IS THE TIME
TO GET STARTED
WITH AJOVY

What is AJOVY[®]?

AJOVY is a prescription medicine used for the preventive treatment of migraine in adults. It is not known if AJOVY is safe and effective in children.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about AJOVY?

Do not use AJOVY if you are allergic to fremanezumab-vfrm or any of the ingredients in AJOVY.

What should I tell my healthcare provider before taking AJOVY?

Before you use AJOVY, tell your healthcare provider if you:

- have high blood pressure.
- have circulation problems in your fingers and toes.

Please see the Important Safety Information throughout and [click here](#) or visit [AJOVY.com](https://www.ajovy.com) to see or print the full [Prescribing Information](#).

AJOVY[®]
(fremanezumab-vfrm)
injection 225 mg/1.5 mL

AJOVY can help reduce monthly migraine days

IN CLINICAL TRIALS, WHEN TAKEN EITHER MONTHLY
OR QUARTERLY OVER A 12-WEEK PERIOD:

50%
OR MORE

About 41% of patients with chronic migraine (vs 18% on placebo) and 48% of patients with episodic migraine (vs 28% on placebo) **had their monthly migraine days reduced by at least 50%**

75%
OR MORE

About 21% of patients with chronic migraine (vs about 10% on placebo) and 27% of patients with episodic migraine (vs 15% on placebo) **had their monthly migraine days reduced by at least 75%**

100%

About 8% of patients with chronic migraine (vs 4% on placebo) and 10% of patients with episodic migraine (vs 5% on placebo) **experienced no migraine days**

Some patients saw results in as early as 1 week.



IMPORTANT SAFETY INFORMATION (CONTINUED)

Before you use AJOVY, tell your healthcare provider if you: (continued)

- are pregnant or plan to become pregnant. It is not known if AJOVY will harm your unborn baby. **Pregnancy Registry:** There is a registry for women who become pregnant during treatment with AJOVY. The purpose of this registry is to collect information about the safety of AJOVY during pregnancy. Contact the registry as soon as you learn that you are pregnant, or ask your doctor to contact the registry for you. You or your doctor can get information and enroll you in the registry by calling 1-833-927-2605 or visiting www.tevamigrainepregnancyregistry.com.

Please see the Important Safety Information throughout and [click here](#) or visit AJOVY.com to see or print the full [Prescribing Information](#).

AJOVY[®]
(fremanezumab-vfrm)
injection 225 mg/1.5 mL

Choose from two different options with AJOVY

AJOVY is the only injectable preventive treatment for migraine that you can take **monthly or quarterly*** with an easy-to-use[†] AJOVY Autoinjector.

MONTHLY

1 injection taken once every month



QUARTERLY

3 injections taken on the same day, once every 3 months



Take a look at these [instructions for how to take AJOVY](#) or visit [AJOVY.com/dosing](https://www.ajovy.com/dosing) and talk to your healthcare provider to decide which dosing option is right for you.

*Monthly dosing is one 225 mg SC injection each month. Quarterly dosing is 675 mg given as three 225 mg SC injections every 3 months.

[†]Two human factor studies tested different people's ability to complete important tasks in order to show how the AJOVY Autoinjector was used in simulated sessions. When asked, "Was the autoinjector easy to use?", 97% in study 1 (N=30) and 98% in study 2 (N=47) answered "Yes."

SC: subcutaneous.

IMPORTANT SAFETY INFORMATION (CONTINUED)

Before you use AJOVY, tell your healthcare provider if you: (continued)

- are breastfeeding or plan to breastfeed. It is not known if AJOVY passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby while using AJOVY.

Please see the Important Safety Information throughout and [click here](#) or visit [AJOVY.com](https://www.ajovy.com) to see or print the full [Prescribing Information](#).

AJOVY[®]
(fremanezumab-vfrm)
injection 225 mg/1.5 mL

Understanding possible side effects with AJOVY

The following side effects were reported by some patients included in AJOVY clinical trials and who were taking AJOVY monthly or quarterly

AJOVY may cause allergic reactions, including itching, rash, and hives that can happen within hours and up to 1 month after taking AJOVY. Adverse reactions reported by $\geq 2\%$ of patients on AJOVY and greater than placebo were injection-site reactions, which were mostly mild to moderate.

2% or less of people stopped taking AJOVY because of side effects.

Less than 1% of people reported constipation or hypertension vs <1% for placebo.

Tell your healthcare provider if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of AJOVY. For more information, ask your healthcare provider or pharmacist. Call your healthcare provider for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.

IMPORTANT SAFETY INFORMATION (CONTINUED)

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Know the medicines you take. Keep a list of your medicines with you to show your healthcare provider and pharmacist when you get a new medicine.

Please see the Important Safety Information throughout and [click here](#) or visit [AJOVY.com](#) to see or print the full [Prescribing Information](#).

AJOVY[®]
(fremanezumab-vfrm)
injection 225 mg/1.5 mL

What to know before taking AJOVY

For subcutaneous* injection only.

Read and follow the Instructions for Use that come with your AJOVY Autoinjector or AJOVY Prefilled Syringe before you start using it and each time you get a refill.

If you still have questions after reviewing, talk to your healthcare provider.

- The AJOVY Autoinjector and AJOVY Prefilled Syringe are for single time (one-time) use only. After use, put the device in an FDA-approved sharps disposal container right away. Do not throw away the sharps disposal container in your household trash.
- Before injecting, let AJOVY sit at room temperature for 30 minutes. Do not warm up using a heat source such as hot water or a microwave.
- After you remove the needle cap from AJOVY, to prevent infection, do not touch the needle.
- Do not pull back on the plunger of the prefilled syringe at any time, as this can break the device.
- Do not inject AJOVY in your veins (intravenously).
- Do not re-use your autoinjector or prefilled syringe as this could cause injury or infection.
- Do not share your device with another person. You may give another person an infection or get an infection from them.
- You may give AJOVY yourself. If you feel uncomfortable, you should not get your first dose of AJOVY until you or your caregiver receive training from a doctor on the right way to use AJOVY.

Storage Conditions:

- Make sure to keep AJOVY in the refrigerator between 36°F to 46°F (2°C to 8°C). Do not freeze.
- Keep AJOVY in the carton it comes in to protect it from light.
- If necessary, AJOVY may be kept in the original carton at room temperature up to 30°C (86°F) for a maximum of 7 days. After removal from the refrigerator, AJOVY must be used within 7 days or discarded. Once stored at room temperature, do not place back in the refrigerator.
- Do not freeze. If AJOVY freezes, throw it away in a sharps disposal container.
- Keep AJOVY out of extreme heat and direct sunlight.
- Do not shake AJOVY.

*Subcutaneous means applied under the skin.

IMPORTANT SAFETY INFORMATION (CONTINUED)

What are the possible side effects of AJOVY?

AJOVY may cause serious side effects, including:

- **Allergic reactions.** Allergic reactions, including itching, rash, and hives, can happen within hours and up to 1 month after receiving AJOVY. Call your healthcare provider or get emergency medical help right away if you have any of the following symptoms of an allergic reaction:
 - swelling of your face, mouth, tongue, or throat
 - trouble breathing

Please see the Important Safety Information throughout and [click here](#) or visit [AJOVY.com](https://www.ajovy.com) to see or print the full [Prescribing Information](#).

AJOVY[®]
(fremanezumab-vfrm)
injection 225 mg/1.5 mL

How to use the AJOVY Autoinjector

Follow these directions whether you take AJOVY monthly or quarterly.

To prepare for your injection, take your autoinjector out of the refrigerator. These are not the complete instructions for using the AJOVY Autoinjector. Read the Patient Information Leaflet located within the full [Prescribing Information](#) before taking AJOVY.

[Watch this video](#) to learn how to use your AJOVY Autoinjector or visit [AJOVY.com/autoinjector/](https://www.ajovy.com/autoinjector/)

STEP 1

Uncap the autoinjector

Pick up the body of the autoinjector with one hand and pull the needle cap straight off with your other hand. Do not shake or twist. Throw cap away immediately.



STEP 2

Place at a 90° angle

Choose an injection area, such as on your stomach, front of thighs, or back of upper arms. Clean the area with an alcohol swab.



STEP 3

Press down and hold the autoinjector until 10 seconds after second click

Press down on the autoinjector. You will hear a click. This means the injection has started and you should see the blue plunger move down. You will hear a second click. Remove the autoinjector 10 seconds after.



IMPORTANT SAFETY INFORMATION (CONTINUED)

What are the possible side effects of AJOVY? (continued)

AJOVY may cause serious side effects, including:

- **High blood pressure.** High blood pressure or worsening of high blood pressure can happen after receiving AJOVY. Contact your healthcare provider if you have an increase in blood pressure.

Please see the Important Safety Information throughout and [click here](#) or visit [AJOVY.com](https://www.ajovy.com) to see or print the full [Prescribing Information](#).

AJOVY[®]
(fremanezumab-vfrm)
injection 225 mg/1.5 mL

How to use the AJOVY prefilled syringe

Follow these directions whether you take AJOVY monthly or quarterly.


To prepare for your injection, take your prefilled syringe out of the refrigerator. These are not the complete instructions for using the AJOVY prefilled syringe. Read the Patient Information Leaflet located within the full [Prescribing Information](#) before taking AJOVY.

[Watch this video](#) to learn how to use your AJOVY prefilled syringe or visit [AJOVY.com/prefilled-syringe/](https://www.ajovy.com/prefilled-syringe/)

STEP 1

Pick up the syringe

Pick up the body of the prefilled syringe with one hand and pull the needle cap straight off with your other hand. Do not shake or twist. Throw cap away immediately.



STEP 2

Select and prepare injection site

Choose an injection area, such as on your stomach, front of thighs, or back of upper arms. Clean the area with an alcohol swab.



IMPORTANT SAFETY INFORMATION (CONTINUED)

What are the possible side effects of AJOVY? (continued)

AJOVY may cause serious side effects, including:

- **Raynaud's phenomenon.** A type of circulation problem can worsen or happen after receiving AJOVY. Raynaud's phenomenon can lead to your fingers or toes feeling numb, cool, or painful, or changing color from pale to blue to red. Contact your healthcare provider if these symptoms occur.

The most common side effects of AJOVY include:

- injection site reactions

Please see the Important Safety Information throughout and [click here](#) or visit [AJOVY.com](https://www.ajovy.com) to see or print the full [Prescribing Information](#).

How to use the AJOVY prefilled syringe (continued)

STEP 3

Insert the needle at a 45° to 90° angle

Use your free hand to gently pinch up at least 1 inch of the injection site. Carefully insert the needle into pinched skin at a 45° to 90° angle.



STEP 4

Slowly push the plunger all the way down

When the needle is all the way into your skin, use your thumb to slowly push the plunger down as far as it will go.



STEP 5

Pull the needle straight out, then apply gauze or cotton ball

After you have injected all of the medicine, pull the needle straight out, then apply gauze or a dry cotton ball on the site for a few seconds.



IMPORTANT SAFETY INFORMATION (CONTINUED)

Tell your healthcare provider if you have any side effect that bothers you or that does not go away.

These are not all the possible side effects of AJOVY. For more information, ask your healthcare provider or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Please see the Important Safety Information throughout and [click here](#) or visit [AJOVY.com](https://www.ajovy.com) to see or print the full [Prescribing Information](#).

AJOVY[®]
(fremanezumab-vfrm)
injection 225 mg/1.5 mL

NOW

**IS THE TIME TO TALK TO YOUR
DOCTOR ABOUT AJOVY**

What is AJOVY?

AJOVY is a prescription medicine used for the preventive treatment of migraine in adults.

It is not known if AJOVY is safe and effective in children.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about AJOVY[®]?

Do not use AJOVY if you are allergic to fremanezumab-vfrm or any of the ingredients in AJOVY.

What should I tell my healthcare provider before taking AJOVY?

Before you use AJOVY, tell your healthcare provider if you:

- have high blood pressure.
- have circulation problems in your fingers and toes.
- are pregnant or plan to become pregnant. It is not known if AJOVY will harm your unborn baby.
Pregnancy Registry: There is a registry for women who become pregnant during treatment with AJOVY. The purpose of this registry is to collect information about the safety of AJOVY during pregnancy. Contact the registry as soon as you learn that you are pregnant, or ask your doctor to contact the registry for you. You or your doctor can get information and enroll you in the registry by calling 1-833-927-2605 or visiting www.tevamigrainepregnancyregistry.com.
- are breastfeeding or plan to breastfeed. It is not known if AJOVY passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby while using AJOVY.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Know the medicines you take. Keep a list of your medicines with you to show your healthcare provider and pharmacist when you get a new medicine.

What are the possible side effects of AJOVY?

AJOVY may cause serious side effects, including:

- **Allergic reactions.** Allergic reactions, including itching, rash, and hives, can happen within hours and up to 1 month after receiving AJOVY. Call your healthcare provider or get emergency medical help right away if you have any of the following symptoms of an allergic reaction:
 - swelling of your face, mouth, tongue, or throat
 - trouble breathing
- **High blood pressure.** High blood pressure or worsening of high blood pressure can happen after receiving AJOVY. Contact your healthcare provider if you have an increase in blood pressure.
- **Raynaud's phenomenon.** A type of circulation problem can worsen or happen after receiving AJOVY. Raynaud's phenomenon can lead to your fingers or toes feeling numb, cool, or painful, or changing color from pale to blue to red. Contact your healthcare provider if these symptoms occur.

The most common side effects of AJOVY include:

- injection site reactions

Tell your healthcare provider if you have any side effect that bothers you or that does not go away.

These are not all the possible side effects of AJOVY. For more information, ask your healthcare provider or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Please see the Important Safety Information throughout and [click here](#) or visit AJOVY.com to see or print the full [Prescribing Information](#).